## **IMMUNIZATION SCHEDULE**





These vaccines are recommended to protect you and to keep your baby safe too!

VACCINES	<b>1-6</b> weeks	<b>7-12</b> weeks	<b>13-19</b> weeks	<b>20-26</b> weeks	<b>27-31</b> weeks	<b>32-36</b> weeks
Flu Vaccine <sup>1</sup> (Influenza)	Yearly, at any point during pregnancy —					
<b>COVID-19 Vaccine</b> (Coronavirus disease 2019)	<ul> <li>✓</li> </ul>	Yearly	/, at any poin	t during preg	nancy ——	
<b>RSV Vaccine</b> <sup>2</sup> Respiratory Syncytial Virus)						Ø
<b>Tdap Vaccine</b> <sup>3</sup> Tetanus, Diphtheria, and Pertussis)						

These shaded boxes show when the vaccine is recommended for pregant people unless your doctor tells you that you cannot safely receive the vaccine.

These vaccines are recommended by the American Academy of Pediatrics: aap.org/immunization

<sup>1</sup> Live attenuated influenza vaccine is not recommended during pregnancy.

<sup>2</sup> 1 dose at 32-36 weeks gestation for babies to be born during RSV season. RSV season is from October to March in the continental United States. Maternal RSV vaccine is given September - January. People who get one dose of RSV vaccine during a pregnancy should not get more doses during future pregnancies. Instead the infant should get nirsevimab.

<sup>3</sup> If Tdap is administered before pregnancy, it should be administered again during pregnancy.

## THESE VACCINES ARE SAFE.

Before being licensed and recommended, each vaccine has been carefully studied by scientific experts at the Food and Drug Administration and CDC.

The CDC continually monitors the safety of all vaccines, which are held to the highest standards.

It is normal and expected to have minor vaccine side effects. Most side effects are mild.

Serious side effects (like severe allergic reaction) are rare.